

Appendix G - Standards for Rangeland Health and Guidelines for Livestock Grazing Management for Public Lands in Oregon and Washington

Introduction

These Standards for Rangeland Health and Guidelines for Livestock Grazing Management for Public Lands in Oregon and Washington were developed in consultation with resource advisory councils and provincial advisory committees, tribes, and others. These standards and guidelines meet the requirements and intent of 43 Code of Federal Regulations, Subpart 4180 (Rangeland Health) and are to be used as presented, in their entirety. These standards and guidelines are intended to provide a clear statement of agency policy and direction for those who use public land for livestock grazing, and for those who are responsible for their management and accountable for their condition. Nothing in this document should be interpreted as an abrogation of Federal trust responsibilities in protection of treaty rights of Indian tribes or any other statutory responsibilities including, but not limited to, the Taylor Grazing Act, Clean Water Act, and Endangered Species Act.

Fundamentals of Rangeland Health

The objectives of the rangeland health regulations referred to above are: “to promote healthy sustainable rangeland ecosystems; to accelerate restoration and improvement of public rangelands to properly functioning conditions ... and to provide for the sustainability of the western livestock industry and communities that are dependent upon productive, healthy public rangelands.”

To help meet these objectives, the regulations on rangeland health identify fundamental principles providing direction to the states, districts, and on-the-ground public land managers and users in the management use of rangeland ecosystems.

A hierarchy, or order, of ecological function and process exists within each ecosystem. The rangeland ecosystem consists of four primary, interactive components; a physical component, a biological component, a social component, and an economic component. This perspective implies that the physical function of an ecosystem supports the biological health, diversity and productivity of that system. In turn, the interaction of the physical and biological components of the ecosystem provides the basic needs of society and supports economic use and potential.

The fundamentals of rangeland health stated in 43 CFR 4180 are:

1. Watersheds are in, or are making significant progress toward, properly functioning physical condition, including their upland, riparian-wetland, and aquatic components; soil and plant conditions support infiltration, soil moisture storage and the release of water that are in balance with climate and landform and maintain or improve water quality and the timing and duration of flow.
2. Ecological processes, including the hydrologic cycle, nutrient cycle and energy flow, are maintained, or there is significant progress toward their attainment, in order to support healthy biotic populations and communities.
3. Water quality complies with State water quality standards and achieves, or is making significant progress toward achieving, established BLM objectives such as meeting wildlife needs.
4. Habitats are, or are making significant progress toward being, restored or maintained for Federal threatened and endangered species, Federal proposed, Category 1 and 2 Federal candidate and other Special Status species.

The fundamentals of rangeland health combine the basic precepts of physical function and biological health elements of law relating to water quality, and plant and animal populations and communities. They provide direction in the development and implementation of the standards for rangeland health.

Standards for Rangeland Health

The standards for rangeland health (standards), based on the above fundamentals, are expressions of the physical and biological condition or degree of function necessary to sustain healthy rangeland ecosystems. Although the focus of these

standards is on domestic livestock grazing on BLM-administered land, on-the-ground decisions must consider the effects and impacts of all issues.

Standards that address the physical components of rangeland ecosystems focus on the roles and interactions of geology and landform, soil, climate, and water as they govern watershed function and soil stability. The biological components addressed in the standards focus on the roles and interactions of plants, animals, and microbes (producers, consumers, and decomposers), and their habitats in the ecosystem. The biological component of rangeland ecosystems is supported by the physical function of the system, and it is recognized that biological activity also influences and supports many of the ecosystem's physical functions.

Guidance contained in 43 CFR 4180 of the regulations directs management toward the maintenance or restoration of the physical function and biological health of rangeland ecosystems. Focusing on the basic ecological health and function of rangelands is expected to provide for the maintenance, enhancement, or creation of future social and economic options.

The standards are based on the ecological potential and capability of each site. In assessing a site's condition or degree of function, it must be understood that the evaluation compares each site to its own potential or capability. Potential and capability are defined as follows:

Potential - The highest level of condition or degree of function a site can attain given no political, social, or economic constraints.

Capability - The highest level of condition or degree of function a site can attain given certain political, social, or economic constraints. For example, these constraints might include riparian areas permanently occupied by a highway or railroad bed that prevent the stream's full access to its original floodplain. If such constraints are removed, the site may be able to move toward its potential.

In designing and implementing management strategies to meet the standards of rangeland health, the potential of the site must be identified, and any constraints recognized, in order that plan goals and objectives are realistic and physically and economically achievable.

Standards and Guidelines in Relation to the Planning Process

The standards apply to the goals of land use plans, activity plans, and project plans (Allotment Management Plans (AMPs)), annual operating plans, habitat management plans, etc.). They establish the physical and biological conditions or degree of function toward which management of publicly-owned rangeland is to be directed. In the development of a plan, direction provided by the standards and the social and economic needs expressed by local communities and individuals are brought together in formulating the goal(s) of that plan.

When the standards and the social and economic goals of the planning participants are woven together in the plan goal(s), the quantifiable, time-specific objective(s) of the plan are then developed. Objectives describe and quantify the desired future conditions to be achieved within a specified timeframe. Each plan objective should address the physical, biological, social, and economic elements identified in the plan goal.

Standards apply to all ecological sites and landforms on public rangelands throughout Oregon and Washington. The standards require site-specific information for full on-the-ground usability. For each standard, a set of indicators is identified for use in tailoring the standards to site-specific situations. These indicators are used for rangeland ecosystem assessments and monitoring, and for developing terms and conditions for permits and leases that achieve the plan goal.

Guidelines for livestock grazing management offer guidance in achieving the plan goal and objectives. The guidelines outline practices, methods, techniques, and considerations used to ensure that progress is achieved in a way, and at a rate, that meets the plan goal and objectives.

Indicators of Rangeland Health

The condition or degree of function of a site, in relation to the standards and its trend toward or away from any standard, is determined through the use of reliable and scientifically sound indicators. The consistent application of such indicators can provide an objective view of the condition and trend of a site when used by trained observers.

For example, the amount and distribution of ground cover can be used to indicate that infiltration at the soil surface can take place as described in the standard relating to upland watershed function. In applying this indicator, the specific levels of plant cover necessary to support infiltration in a particular soil should be identified using currently available information from reference areas, if they exist; from technical sources like soil survey reports, ecological site inventories, and ecological site descriptions, or from other existing reference materials. Reference areas are land that best represent the potential of a specific ecological site in both physical function and biological health. In many instances, potential reference areas are identified in ecological site descriptions and are referred to a “type location.” In the absence of suitable reference areas, the selection of indicators to be used in measuring or judging condition or function should be made by an interdisciplinary team of experienced professionals and other trained individuals.

Not all indicators identified for each standard are expected to be employed in every situation. Criteria for selecting appropriate indicators and methods of measurement and observation include, but are not limited to, 1) the relationship between the attribute(s) being measured or observed and the desired outcome; 2) the relationship between the activity (e.g., livestock grazing) and the attribute(s) being measured or observed, and 3) funds and workforce available to conduct the measurements or observations.

Assessment and Monitoring

The standards are the basis for assessing and monitoring rangeland condition and trend. Carrying out well-designed assessment and monitoring is critical to restoring or maintaining healthy rangelands and determining trends and conditions.

Assessments are a cursory form of evaluation based on the standards that can be used at different landscape scales. Assessments, conducted by qualified interdisciplinary teams (which may include, but are not limited to, physical, biological, and social specialists and interagency personnel) with participation from permittees and other interested parties, are appropriate at the watershed and subwatershed level, at the allotment and pasture levels, and on individual ecological sites or groups of sites. Assessments identify the condition or degree of function within the rangeland ecosystem and indicate resource problems and issues that should be monitored or studied in more detail. The results of the assessments are a valuable tool for managers in assigning priorities within an administrative area and the subsequent allocation of personnel, money, and time in resource monitoring and treatment. The results of assessments may also be used in making management decisions where an obvious problem exists.

Monitoring, which is the well-documented and orderly collection, analysis, and interpretation of resource data, serves as the basis for determining trends in the condition or degree of function of rangeland resources and for making management decisions. Monitoring should be designed and carried out to identify trends in resource conditions, to point out resource problems, to help indicate the cause of such problems, to point out solutions, and/or to contribute to adaptive management decisions. In cases where monitoring data do not exist, professional judgement, supported by interdisciplinary team recommendation, may be relied upon by the authorized officer in order to take necessary action. Review and evaluation of new information must be an ongoing activity.

To be effective, monitoring must be consistent over time, throughout administrative areas, and in the methods of measurement and observation of selected indicators. Those doing the monitoring must have the knowledge and skill required by the level or intensity of the monitoring being done, as well as the experience to properly interpret the results. Technical support for training must be made available.

Measurability

It is recognized that not every area will immediately meet the standards and that it will sometimes be a long-term process to restore some rangelands to properly functioning condition. It is intended that in cases where standards are not being met, measurable progress should be made toward achieving those standards, and significant progress should be made toward fulfilling the fundamentals of rangeland health. Measurability is defined on a case-specific basis based upon the stated planning objectives (e.g., quantifiable, time-specific), taking into account economic and social goals along with the biological and ecological capability of the area. To the extent that a rate of recovery conforms with the planning objectives, the area is allowed the time to meet the standard under the selected management regime.

Implementation

The material contained in this document will be incorporated into existing land use plans and used in the development of new land use plans. According to 43 CFR 4130.3-1, permits and leases shall incorporate terms and conditions that

ensure conformance with 43 CFR 4180. Terms and conditions of existing permits and leases will be modified to reflect standards and guidelines at the earliest possible date, with priority for modification being at the discretion of the authorized officer. Terms and conditions of new permits and leases will reflect standards and guidelines in their development.

Indicators identified in this document will serve as a focus of interpretation of existing monitoring data and will provide the basis of design for monitoring and assessment techniques, and in the development of monitoring and assessment plans.

The authorized officer shall take appropriate action as soon as practicable, but not later than the start of the next grazing year, upon determining through assessment or monitoring by experienced professionals and interdisciplinary teams that a standard is not being achieved and that livestock are a significant contributing factor to the failure to achieve the standards and conform with the guidelines.

Standards for Rangeland Health

Standard 1: Watershed Function - Uplands

Upland soils exhibit infiltration and permeability rates, moisture storage, and stability that are appropriate to soil, climate, and landform.

Rationale and Intent:

This standard focuses on the basic physical functions of upland soils that support plant growth, the maintenance or development of plant populations and communities, and promote dependable flows of quality water from the watershed.

To achieve and sustain rangeland health, watersheds must function properly. Watersheds consist of three principle components; the uplands, riparia/wetland areas, and the aquatic zone. This standard addresses the upland component of the watershed. When functioning properly, within its potential, a watershed captures, stores, and safely releases the moisture associated with normal precipitation events (equal to or less than the 25-year, 5-hour event) that falls within its boundaries. Uplands make up the largest part of the watershed and are where most of the moisture is received during precipitation events is captured and stored.

While all watersheds consist of similar components and processes, each is unique in its individual makeup. Each watershed displays its own pattern of landform and soil, its unique climate and weather patterns, and its own history of use and current condition. In directing management toward achieving this standard, it is essential to treat each unit of the landscape (soil, ecological site, and watershed) according to its own capability and how it fits with both smaller and larger units of the landscape.

A set of potential indicators has been identified for which site-specific criteria will be used to determine if this standard is being met. The appropriate indicators to be used in determining attainment of the standard should be drawn from the following list.

Potential Indicators:

Protection of the soil surface from raindrop impact; detention of overland flow; maintenance of infiltration and permeability, and protection of the soil surface from erosion, consistent with the potential/capability of the site, as evidenced by the:

- amount and distribution of plant cover (including forest canopy cover);
- amount and distribution of plant litter;
- accumulation/incorporation of organic matter;
- amount and distribution of bare ground;
- amount and distribution of rock, stone, and gravel;

- plant composition and community structure;
- thickness and continuity of the “A” horizon;
- character of microrelief;
- presence and integrity of biotic crusts;
- root occupancy of the soil profile;
- biological activity (plant, animal, and insect); and
- absence of accelerated erosion and overland flow.

Soil and plant conditions promote moisture storage as evidenced by:

- amount and distribution of plant cover (including canopy cover);
- amount and distribution of plant litter;
- plant composition and community structure; and
- accumulation/incorporation of organic matter.

Standard 2: Watershed Function - Riparian/Wetland Areas

Riparian/wetland areas are in properly functioning physical condition appropriate to soil, climate and landform.

Rational and Intent:

Riparian/wetland areas are grouped into two major categories: 1) lentic, or standing water systems such as lakes, ponds, seeps, bogs, and meadows; and 2) lotic, or moving water systems such as rivers, streams, and springs. Wetlands are areas that are inundated or saturated by surface or ground water at a frequency and duration to support, and which under normal circumstances do support, a prevalence of vegetation typically adapted to life in saturated soil conditions. Riparian areas commonly occupy the transition zone between the upland and surface water bodies (the aquatic zone) or permanently saturated wetlands.

Properly functioning condition of riparian and wetland areas describes the degree of physical function of these components of the watershed. Their functionality is important to water quality in the capture and retention of sediment and debris, the detention and detoxification of pollutants, and in moderating seasonal extremes of water temperature. Properly functioning riparian areas and wetlands enhance the timing and duration of streamflow through dissipation of flood energy, improved bank storage, and ground water recharge. Properly functioning condition should not be confused with the desired plant community or the desired future condition since, in most cases, it is the precursor to these levels of resource condition and is required for their attainment.

A set of indicators has been identified for which site-specific criteria will be used to determine if this standard is being met. The criteria are based upon the potential (or upon the capability where potential cannot be achieved) of individual sites or landforms.

Potential Indicators:

Hydrologic, vegetation, and erosional/depositional processes interact in supporting physical function, consistent with the potential or capability of the site, as evidenced by:

- frequency of floodplain/wetland inundation;
- plant composition, age class distribution, and community structure;

- root mass;
- point bars revegetating;
- streambank/shoreline stability;
- riparian area width;
- sediment deposition;
- active/stable beaver dams;
- coarse/large woody debris;
- upland watershed conditions;
- water table fluctuation.

Stream channel characteristics are appropriate for landscape position as evidenced by:

- channel width/depth ratio;
- channel sinuosity;
- gradient;
- rocks and coarse and/or large woody debris;
- overhanging banks;
- pool/riffle ratio;
- pool size and frequency; and
- stream embeddedness.

Standard 3: Ecological Processes

Healthy, productive, and diverse plant and animal populations and communities appropriate to soil, climate, and landform are supported by ecological processes of nutrient cycling, energy flow, and the hydrologic cycle.

Rationale and Intent:

This standard addresses the ecological processes of energy flow and nutrient cycling as influenced by existing and desired plant and animal communities without establishing the kinds, amounts, or proportions of plant and animal community compositions. While emphasis may be on native species, an ecological site may be capable of supporting a number of different native and introduced plant and animal populations and communities while meeting this standard. This standard also addresses the hydrologic cycle which is essential for plant growth and appropriate levels of energy flow and nutrient cycling. Standards 1 and 2 address the watershed aspects of the hydrologic cycle.

With a few exceptions, all life on earth is supported by the energy supplied by the sun and captured by plants in the process of photosynthesis. This energy enters the food chain when plants are consumed by insects and herbivores and passes upward through the food chain to the carnivores. Eventually, the energy reaches the decomposers and is released as the thermal output of decomposition or through oxidation.

The ability of plants to capture sunlight energy, to grow and develop, to play a role in soil development and watershed function, to provide habitat for wildlife, and to support economic uses depends on the availability of nutrients and moisture. Nutrients necessary for plant growth are made available to plants through the decomposition and metabolization of organic matter by insects, bacteria and fungi, the weathering of rocks, and extraction from the

atmosphere. Nutrients are transported through the soil by plant uptake, leaching, and by rodent, insect, and microbial activity. They follow cyclical patterns as they are used and reused by living organisms.

The ability of rangelands to supply resources and satisfy social and economic needs depends on the buildup and cycling of nutrients over time. Interrupting or slowing nutrient cycling can lead to site degradation, as this land becomes increasingly deficient in the nutrients plants require.

Some plant communities, because of past use, frequent fire or other histories of extreme or continued disturbance, are incapable of meeting this standard. For example, shallow-rooted winter-annual grasses that completely dominate some sites do not fully occupy the potential rooting depth of some soils, thereby reducing nutrient cycling well below optimum levels. In addition, these plants have a relatively short growth period and thus capture less sunlight than more diverse plant communities. Plant communities like those cited in this example are considered to have crossed the threshold of recovery and often require great expense to be recovered. The cost of recovery must be weighed against the site's potential ecological/economic value in establishing treatment priorities.

The role of indicators has been identified for which site-specific criteria will be used to determine if this standard is being met.

Potential Indicators:

Photosynthesis is effectively occurring throughout the potential growing season, consistent with the potential/capability of the site, as evidenced by plant composition and community structure.

Nutrient cycling is occurring effectively, consistent with the potential/capability of the site, as evidenced by:

- plant composition and community structure;
- accumulation, distribution, incorporation of plant litter and organic matter into the soil;
- animal community structure and composition;
- root occupancy in the soil profile; and
- biological activity including plant growth, herbivory, and rodent, insect, and microbial activity.

Standard 4: Water Quality

Surface water and ground water quality, influenced by agency actions, complies with State water quality standards.

Rationale and Intent:

The quality of the water yielded by a watershed is determined by the physical and chemical properties of the geology and soils unique to the watershed, the prevailing climate and weather patterns, current resource conditions, the uses to which the land is put, and the quality of the management of the uses. Standards 1, 2, and 3 contribute to attaining this standard.

States are legally required to establish water quality standards and Federal land management agencies are to comply with those standards. In mixed ownership watersheds, agencies, like any other landowners, have limited influence on the quality of the water yielded by the watershed. The actions taken by the agency will contribute to meeting State water quality standards during the period that water crosses agency administered holdings.

Potential Indicators:

Water quality meets applicable water quality standards as evidenced by:

- water temperature;
- dissolved oxygen;

- fecal coliform;
- turbidity;
- pH;
- populations of aquatic organisms; and
- effects on beneficial uses (e.g., effects on management activities on beneficial uses as defined under the Clean Water Act and State implementing regulations).

Standard 5: Native, Threatened and Endangered, and Locally Important Species

Habitats support healthy, productive, and diverse populations and communities of native plants and animals (including Special Status species and species of local importance) appropriate to soil, climate, and landform.

Rationale and Intent:

Federal agencies are mandated to protect threatened and endangered species and will take appropriate action to avoid the listing of any species. This standard focuses on retaining and restoring native plant and animal (including fish) species, populations, and communities (including threatened, endangered and other Special Status species and species of local importance). In meeting the standard, native plant communities and animal habitats would be spatially distributed across the landscape with a density and frequency of species suitable to ensure reproductive capability and sustainability. Plant populations and communities would exhibit a range of age classes necessary to sustain recruitment and mortality fluctuations.

Potential Indicators:

Essential habitat elements for species, populations, and communities are present and available, consistent with the potential/capability of the landscape, as evidenced by:

- plant community composition, age class distribution, productivity;
- animal community composition, productivity;
- habitat elements;
- spatial distribution of habitat;
- habitat connectivity; and
- population stability/resilience.

Guidelines for Livestock Grazing Management

Guidelines for livestock grazing management offer guidance in achieving plan goals, meeting standards for rangeland health, and fulfilling the fundamentals of rangeland health. Guidelines are applied in accordance with the capabilities of the resource in consultation, cooperation, and coordination with permittees/lessees and the interested public. Guidelines enable managers to adjust grazing management on public land to meet current and anticipated climatic and biological conditions.

General Guidelines

1. Involve diverse interests in rangeland assessment, planning, and monitoring.
2. Assessment and monitoring are essential to the management of rangelands, especially in areas where resource problems exist or issues arise. Monitoring should proceed using a qualitative method of assessment to identify

critical, site-specific problems or issues using interdisciplinary teams of specialists, managers, and knowledgeable land users.

Once identified, critical, site-specific problems or issues should be targeted for more intensive, quantitative monitoring or investigation. Priority for monitoring and treatment should be given to those areas that are ecologically at-risk where benefits can be maximized given existing budgets and other resources.

Livestock Grazing Management

1. The season, timing, frequency, duration, and intensity of livestock grazing use should be based on the physical and biological characteristics of the site and the management unit in order to:
 - a. Provide adequate cover (live plants, plant litter, and residue) to promote infiltration, conserve soil moisture, and to maintain soil stability in upland areas;
 - b. Provide adequate cover and plant community structure to promote streambank stability, debris and sediment capture, and floodwater energy dissipation in riparian areas;
 - c. Promote soil surface conditions that support infiltration;
 - d. Avoid subsurface soil compaction that retards the movement of water in the soil profile;
 - e. Help prevent the increase and spread of noxious weeds;
 - f. Maintain or restore diverse plant populations and communities that fully occupy the potential rooting volume of the soil;
 - g. Maintain or restore plant communities to promote photosynthesis throughout the potential growing season;
 - h. Promote soil and site conditions that provide the opportunity for the establishment of desirable plants;
 - i. Protect or restore water quality; and
 - j. Provide for the life cycle requirements, and maintain or restore the habitat elements of native (including threatened and endangered, Special Status, and locally important species) and desired plants and animals.
2. Grazing management plans should be tailored to site-specific conditions and plan objectives. Livestock grazing should be coordinated with the timing of precipitation, plant growth, and plant form. Soil moisture, plant growth stage, and the timing of peak streamflows are key factors in determining when to graze. Response to different grazing strategies varies with differing ecological sites.
3. Grazing management systems should consider nutritional and herd health requirements of the livestock.
4. Integrate grazing management systems into the year-round management strategy and resources of the permittee(s) or lessee(s). Consider the use of collaborative approaches (e.g., coordinated resource management, work groups) in this integration.
5. Consider competition for forage and browse among livestock, big game animals, and wild horses in designing and implementing a grazing plan.
6. Provide periodic rest from grazing for rangeland vegetation during critical growth periods to promote plant vigor, reproduction, and productivity.
7. Range improvement practices should be prioritized to promote rehabilitation and resolve grazing concerns on transitory grazing land.

8. Consider the potential for conflict between grazing use on public land and adjoining land uses in the design and implementation of a grazing management plan.

Facilitating the Management of Livestock Grazing

1. The use of practices to facilitate the implementation of grazing systems should consider the kind and class of animals managed, indigenous wildlife, wild horses, the terrain, and the availability of water. Practices such as fencing, herding, water development, and the placement of salt and supplements (where authorized) are used where appropriate to:
 - a. Promote livestock distribution;
 - b. Encourage a uniform level of proper grazing use throughout the grazing unit;
 - c. Avoid unwanted or damaging concentrations of livestock on streambanks, in riparian areas, and other sensitive areas such as highly erodible soils, unique wildlife habitats, and plant communities; and
 - d. Protect water quality.
2. Roads and trails used to facilitate livestock grazing are constructed and maintained in a manner that minimizes the effects on landscape hydrology; concentration of overland flow, erosion, and sediment transport are prevented; and subsurface flows are retained.

Accelerating Rangeland Recovery

1. Upland treatments that alter the vegetation composition of a site, such as prescribed burning, juniper management, and seedings or plantings must be based on the potential of the site and should:
 - a. Retain or promote infiltration, permeability, and soil moisture storage;
 - b. Contribute to nutrient cycling and energy flow;
 - c. Protect water quality;
 - d. Help prevent the increase and spread of noxious weeds;
 - e. Contribute to the diversity of plant communities, and plant community composition and structure;
 - f. Support the conservation of threatened and endangered, other Special Status species, and species of local importance; and
 - g. Be followed up with grazing management and other treatments that extend the life of the treatment and address the cause of the original treatment need.
2. Seedings and plantings of nonnative vegetation should only be used in those cases where native species are not available in sufficient quantities; where native species are incapable of maintaining or achieving the standards; or where nonnative species are essential to the functional integrity of the site.
3. Structural and vegetation treatments and animal introductions in riparian and wetland areas must be compatible with the capability of the site, including the system's hydrologic regime, and contribute to the maintenance or restoration of properly functioning condition.

Rangelands Glossary

Appropriate action - implementing actions pursuant to subparts 4110, 4120, 4130, and 4160 of the regulations that will result in significant progress toward fulfillment of the standards and significant progress toward conformance with the guidelines. (See Significant progress")

Assessment - a form of evaluation based on the standards of rangeland health, conducted by an interdisciplinary team at the appropriate landscape scale (pasture, allotment, subwatershed, watershed, etc.) to determine conditions relative to standards.

Compaction layer - a layer within the soil profile in which the soil particles have been rearranged to decrease void space, thereby increasing soil bulk density and often reducing permeability.

Crust, Abiotic - (physical crust) a surface layer on soils, ranging in thickness from a few millimeters to a few centimeters, that is much more compact, hard, and brittle when dry, than the material immediately beneath it.

Crust, Biotic - (microbiotic or cryptogamic crust) a layer of living organisms (mosses, lichens, liverworts, algae, fungi, bacteria, and/or cyanobacteria) occurring on, or near the soil surface.

Degree of function - a level of physical function relative to properly functioning condition commonly expressed as: properly functioning, functioning-at-risk, or nonfunctional.

Diversity - the aggregate of species assemblages (communities), individual species, and the genetic variation within species and the processes by which these components interact within and among themselves. The elements of diversity are: 1) community diversity (habitat, ecosystem); 2) species diversity; and 3) genetic diversity within a species; all three of which change over time.

Energy flow - the processes in which solar energy is converted to chemical energy through photosynthesis and passed through the food chain until it is eventually dispersed through respiration and decomposition.

Ground water - water in the ground that is in the zone of saturation; water in the ground that exists at, or below the water table.

Guideline - practices, methods, techniques, and considerations used to ensure that progress is made in a way and at a rate that achieves the standard(s).

Gully - a channel resulting from erosion and caused by the concentrated but intermittent flow of water usually during and immediately following heavy rains.

Hydrologic cycle - the process in which water enters the atmosphere through evaporation, transpiration, or sublimation from the oceans, other surface water bodies, or from the land and vegetation, and through condensation and precipitation returns to the earth's surface. The precipitation then occurring as overland flow, streamflow, or percolating underground flow to the oceans or other surface water bodies or to other sites of envirotranspiration and recirculation to the atmosphere.

Indicators - parameters of ecosystem function that are observed, assessed, measured, or monitored to directly or indirectly determine attainment of a standard(s).

Infiltration - the downward entry of water into the soil.

Infiltration rate - the rate at which water enters the soil.

Nutrient cycling - the movement of essential elements and inorganic compounds between the reservoir pool (soil, for example) and the cycling pool (organisms) in the rapid exchange (e.g., moving back and forth) between organisms and their immediate environment.

Organic matter - plant and animal residues accumulated or deposited at the soil surface; the organic fraction of the soil that includes plant and animal residues at various stages of decomposition; cells and tissues of soil organisms, and the substances synthesized by the soil population.

Permeability - the ease with which gases, liquids, or plant roots penetrate or pass through a bulk mass of soil or a layer of soil.

Properly functioning condition - *Riparian/wetland*: adequate vegetation, landform, or large (coarse) woody debris is present to dissipate stream energy associated with high waterflows, thereby reducing erosion and improving water

quality; filter sediment, capture bedload, and aid in floodplain development; improve floodwater retention and ground water recharge; develop root masses that stabilize streambanks against cutting action; develop diverse channel and ponding characteristics to provide the habitat and water depth, duration, and temperature necessary for fish production, waterfowl breeding, and other uses; and support greater biodiversity. The result of interaction among geology, soil, water, and vegetation. *Uplands*: soil and plant conditions support the physical processes of infiltration and moisture storage and promote soil stability (as appropriate to site potential); includes the production of plant cover and the accumulation of plant residue that protect the soil surface from raindrop impact, moderate soil temperature in minimizing frozen soil conditions (frequency, depth, and duration), and the loss of soil moisture to evaporation; root growth and development in the support of permeability and soil aeration. The result of interaction among geology, climate, landform, soil, and organisms.

Proper grazing use - grazing that, through the control of timing, frequency, intensity, and duration of use, meets the physiological needs of the desirable vegetation, provides for the establishment of desirable plants, and is in accord with the physical function and stability of soil and landform (properly functioning condition).

Reference area - site that, because of their condition and degree of function, represent the ecological potential or capability of similar sites in an area or region (ecological province); serve as a benchmark in determining the ecological potential of sites with similar soil, climatic, and landscape characteristics.

Rill - a small, intermittent water course with steep sides; usually only a few inches deep.

Riparian area - a form of wetland transition between permanently saturated wetlands and upland areas. These areas exhibit vegetation or physical characteristics reflective of permanent surface or subsurface water influence. Land along, adjacent to, or contiguous with perennially and intermittently flowing rivers and stream, glacial potholes, and shores of lakes and reservoirs with stable water levels are typical riparian areas. Excluded are such sites as ephemeral streams or washes that do not exhibit the presence of vegetation dependent upon free water in the soil. Includes, but is not limited to, jurisdictional wetlands.

Significant progress - when used in reference to achieving a standard: (actions), the necessary land treatments, practices, and/or changes to management have been applied or are in effect; (rate), a rate of progress that is consistent with the anticipated recovery rate described in plan objectives, with due recognition of the effects of climatic extremes (drought, flooding, etc.), fire, and other unforeseen naturally occurring events or disturbances. Monitoring reference areas that are ungrazed and properly grazed may provide evidence of appropriate recovery rates. (See Proper Grazing Use)

Soil density - (bulk density) - the mass of dry soil per unit bulk volume.

Soil moisture - water contained in the soil; commonly used to describe water in the soil above the water table.

Special Status species - species proposed for listing, officially listed (threatened/endorsed), or candidate for listing as threatened or endorsed by the Secretary of the Interior under the provisions of the Endangered Species Act; those listed or proposes for listing by the State in a category implying potential endorsement or extinction; those designated by each BLM State Director as sensitive.

Species of local importance - species of significant importance to American Indian populations (e.g., medicinal and food plants).

Standard - an expression of the physical and biological condition or degree of function necessary to sustain healthy rangeland ecosystems.

Uplands - land that exists above the riparian/wetland area, or active floodplains of rivers and streams; those lands not influenced by the water table or by free or unbound water; commonly represented by toe slopes, alluvial fans, and side slopes, shoulders, and ridges of mountains and hills.

Watershed - an area of land that contributes to the surface flow of water past a given point. The watershed dimensions are determined by the point past, or through which, runoff flows.

Watershed function - the principal functions of a watershed include the capture of moisture contributed by precipitation; the storage of moisture within the soil profile, and the release of moisture through the subsurface flow, deep percolation to ground water, evaporation from the soil, and transpiration by live vegetation.

Wetland - areas that are inundated or saturated by surface or ground water at a frequency and duration sufficient to support, and which under normal circumstances do support, a prevalence of vegetation typically adapted for life in saturated soil conditions.

This Page Intentionally Left Blank