

Appendix E

South Steens Allotment Monitoring Data

Site	Year	Type of Use	Plant Species	Utilization Level
1	1996	Cattle and Horses	Idaho Fescue	38% Light
1	1997	Horses	Idaho Fescue	65% Heavy
2	1997	Horses	Physical Damage Riparian	40% Light
2	1995	Horses	Physical Damage Meadows	NA
3	1997	Horses	Riparian	70% Heavy
3	1996	Cattle and Horses	Riparian	69% Heavy
4A	1997	Horses	Idaho Fescue	45% Moderate
4B	1997	Horses	Riparian	90% Severe
4A	1996	Cattle and Horses	Idaho Fescue	45% Moderate
4B	1996	Cattle and Horses	Riparian	90% Severe
5	1997	Cattle and Horses	Riparian	16% Slight
5	1996	Cattle and Horses	Riparian	48% Moderate
6	1997	Horses	Idaho Fescue	69% Heavy
6	1996	Cattle and Horses	Idaho Fescue	49% Moderate
7A	1997	Horses	Thurber's needlegrass	50% Moderate
7A	1996	Cattle and Horses	Thurber's needlegrass	51% Moderate
7B	1997	Horses	Riparian	90% Severe
7B	1996	Cattle and Horses	Riparian	90% Severe
8	1997	Horses	Riparian	30% Light
8	1996	Cattle and Horses	Riparian	35% Light
9	1997	Cattle and Horses	Riparian	37% Light
9	1996	Cattle and Horses	Riparian	49% Moderate
10	1997	Horses	Riparian	70% Heavy